

Cost of Disease Modifying Therapies for Multiple Sclerosis: Is Front-Loading the Answer?

Lisa Batcheller, David Baker

Barts and the London School of Medicine and Dentistry

Queen Mary University of London, United Kingdom

Correspondence: Lisa Batcheller, Blizard Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, 4 Newark Street, London E1 2AT. United Kingdom.
Tel: +44 (0)20 7882 2485, Fax 020 7882 2180, Email L.Batcheller@smd12.qmul.ac.uk.

Conflicts of Interest: LB has none to declare. DB. Non-considered relevant but is a shareholder of Canbex therapeutics and has received honoraria from in the past 3 years: Japan Tobacco; Merck, Roche; Sanofi-Genzyme.

Abstract

There are now over a dozen disease modifying therapies (DMTs) available to treat multiple sclerosis (MS). They vary in efficacy and safety as well as in cost.

The literature on the cost effectiveness of these is often confusing and contradictory. There is a lack of quality evidence enabling the comparison of different DMTs. There are scarce randomised controlled trials which look at one DMT compared with another that is not IFN or GA. There is also a lack of systematic reviews comparing the efficacy and safety of different DMTs. This makes it difficult to perform good quality cost-effectiveness analyses (CEAs). Furthermore, CEAs in and of themselves are difficult to interpret or compare due to the variation in methods and cost estimations as well as the use of outcome measures which cannot be proven over a reasonable timeframe.

This review looks at the different DMTs available for MS and attempts to draw some conclusions on their cost-effectiveness. It also considers the costs and benefits of front loading the cost of treatment for MS by using more expensive and effective treatment earlier on.

Multiple Sclerosis

Multiple sclerosis (MS) is a chronic autoimmune inflammatory condition of the central nervous system (CNS). It is a pathophysiological process of inflammation, demyelination and neurodegeneration (1). It can cause severe disability as function of the CNS deteriorates (1). Until the 1990s, treatment for MS consisted only of symptomatic control, usually a short course of high dose steroids during exacerbations of the disease (2). MS attacks would lead to progressively worsening disability and early death (3). Since then, a multitude of disease modifying therapies (DMTs) have been developed that can slow the course of the disease. These drugs (Table 1) are high-cost and not without their own risks, but are now widely used to treat relapsing-remitting forms of MS (RRMS).

Diagnosis and Clinical Course

MS is diagnosed when clinical features and investigations indicate there is focal demyelination occurring in more than one location and on more than one occasion, such that there is dissemination in space and dissemination in time (4). Diagnosis of MS often requires the exclusion of other possible diagnoses as presentation is varied (5). Clinical features of MS are many and include motor, sensory and autonomic disturbances (1). Most commonly, MS presents with an acute attack of demyelination resulting in neurological dysfunction (1). In 80% of cases, this will resolve before subsequent attacks occur (1). This is the relapsing-remitting course of the disease. Over time, recovery between episodes diminishes and symptoms become progressively worse without recovery and this is termed secondary progressive MS (SPMS). About 15-20% of people with MS will have a progressive course from onset, known as primary progressive MS (PPMS) (1).

Pathogenesis

There are several proposed mechanisms by which CNS autoimmunity may be initiated in MS. Molecular mimicry may initiate the process by which autoreactive T cells are stimulated by autoantigens (6). By another proposed mechanism, antigens, such as glial or neural proteins may drain into cervical lymph nodes via the glymphatic drainage system (7, 8). Once in the cervical lymph nodes, these antigens are presented to autoreactive T cells by antigen presenting cells (APCs) (6). There seems to be a failure of regulatory T cells to suppress autoreactive T cells upon activation by an autoantigen (9). These autoreactive T cells are able to extravasate into the CNS and they are then re-activated locally, causing the release of cytokines which promote recruitment of microglia and directly cause damage to myelin and axons (10). Autoreactive T cells will also activate B cells and a humoral response will further add to inflammatory cell recruitment and contribute to damage to myelin and axons (11). Alternatively, it may be that memory B cells are the major driving force behind the inflammatory phase in MS and present antigens to T cells or directly damage the CNS via toxic molecules or stimulate the innate immune response (12, 13). Nearly all DMTs seem to deplete memory B cells, and the degree of depletion can be correlated to the efficacy of the drug (12, 13).

Impact

MS carries a huge burden for those afflicted with the disease and those close to them. People with MS experience a decreased quality of life, and life in the most severe stages of MS can be considered to be worse than death (14, 15). Not only do people with MS suffer from physical disability, cognitive impairment can also occur from the earliest stages of the disease onwards (16). People with MS are

more likely to suffer from anxiety and depression than the general population, and they are also more likely to commit suicide (17, 18). Furthermore, people with MS are 40% more likely to get a divorce compared to matched controls that did not have MS (19).

MS has an effect on the ability of people to work full-time and in most cases eventually leads to unemployment. A cost-of-illness study carried out in 2017 looking at data from 16,808 people with MS across 16 European countries found that overall average unemployment was 68% (20).

Relapses can be disabling for people with MS and costly for healthcare systems. In the 2017 European study, 13% of people had experienced a relapse and 7.5% had been admitted to hospital as an inpatient in the last 3 months (20). Overall, 57% of people were using DMTs, with usage varying by country from 26-79% (20).

The 2017 study found that in the UK the total annual cost to society ranged from £11,400 (£12,800) for mild disease (EDSS 0-3) to £22,700 (£25,400) for the moderate disease group (EDSS 4-6.5) and £36,500 (£40,900) for the severe disease group (EDSS 7-9) (21). Overall, the average cost by disease severity across Europe was €22,800, €37,100 and €57,500 for mild, moderate and severe disease (20). For less severe disease, the majority of direct costs were due to health care usage, whereas informal care costs accounted for most of the direct costs in more severe disease (20). Production losses accounted for 39%, 45% and 33% of total costs for mild, moderate and severe disease groups respectively (20).

Table 1: Characteristics of Disease Modifying Therapies for Multiple Sclerosis								
DMT	Dose	Pivotal trial efficacy	MoA	Side effects	Monitoring	Annual Cost (UK)*	First year Cost (US) (Year 1 WAC)	EMA Licensing
IFNb-1a (Rebif s.c. injection) (Avonex i.m. injection)	Weekly i.m. injection (30µg) or thrice weekly s.c. injection (20 or 44µg)	s.c.: PRISMS – 27% and 33% reduction in ARR compared to placebo for the 22µg and 44µg dose groups respectively, Increased time to SAD compared to placebo (22) i.m.: 18% reduction in ARR compared to placebo, 37% reduction in SAD vs placebo (23)	The are several proposed mechanisms of action for beta-interferons, including inhibition of immune cell proliferation, inhibition of B and T cell activation (24, 25), increasing anti-inflammatory cytokines (26) , increasing apoptosis of autoreactive T cells (27) and limiting T cell-endothelial adhesion (28, 29).	Injection site reactions, hepatic impairment, depression, flu-like symptoms. Rarer side effects include anaemia, thrombocytopenia, lymphopenia and thrombotic microangiopathy (30, 31). There is an association with capillary leak syndrome in undiagnosed monoclonal gammopathy, which is why the Association of British Neurologists recommend that protein electrophoresis should be carried out prior to commencing therapy (32, 33)	FBC, LFT, renal function, blood pressure at baseline and every 3-6 months Protein electrophoresis at baseline (33, 34)	Rebif: 20/44µg £9,088 /£12,068 Avonex: £9,061	Rebif: \$86,416 Avonex: \$81,965	RRMS
IFNb-1b (Betaseron or Extavia)	s.c. injection every	34% reduction in ARR compared to placebo, no				Betaseron And Extavia: £7,264	Betaseron: \$86,659 Extavia: \$72,359	

Pre-submission

	other day (250µg)	significant effect on SAD (35)						
Pegylated IFNβ-1a (Plegidry)	s.c. injection every two weeks (125µg)	ADVANCE – 35% reduction in ARR and a 39% reduction in 3- month SAD compared to placebo (36)		Flu like symptoms, headache, injection site reactions, raised liver enzymes and depression are common side effects Rarer side effects include hypersensitivity reactions, seizures, injection site reactions, renal damage (37)	Regular monitoring of FBC, LFT, TFT, blood pressure and renal function (37)	£8,500	\$81,956	RRMS
Glatiramer Acetate (GA) (Copaxone or Glatopa (generic))	Daily s.c. injection (20mg) Or thrice weekly s.c. injection (40mg)	29% reduction in ARR vs placebo, no significant difference in SAD (38)	GA promotes apoptosis of autoreactive cells by increasing levels of a strongly suppressive subset of regulatory T cells (PD1- Treg cells) (27). Furthermore GA is thought to interfere with antigen presentation and T cell activation, as well as increasing anti- inflammatory cytokines (39, 40).	Injection site reactions, Symptoms mimicking cardiac ischaemia – chest tightness, shortness of breath, palpitations, anxiety (31)	No additional monitoring required	£6,700	20mg: \$86,554 40mg: \$76,024 20mg Glatopa (generic): \$63,193	RRMS
Fingolimod (Gilenya)	Daily oral tablet (0.5mg)	FREEDOMS II – 48% reduction in ARR compared to placebo, no significant reduction in SAD (41) TRANSFORMS – 52% reduction in ARR compared to i.m. IFNβ1a, no significant difference in SAD (42)	Fingolimod is a sphingoside-1- phosphate (S1P) receptor antagonist. S1P receptors are expressed in a variety of cells including lymphocytes, neurons and astroglia. Binding of S1P to S1P receptors mediates lymphocyte migration out of lymph glands, and this process is disrupted by fingolimod as it is an S1P analogue. This mainly affects central memory T cells and naïve T cells as these are more likely to re- circulate through lymph glands (43).	First dose bradycardia, First and second degree atrioventricular block, Macular oedema, Lymphopenia, Infections, Elevated liver enzymes, Hypertension (34, 41) Rarely PML (44)	MRI, ECG, VZV IgG serology, pregnancy test, ophthalmology testing, FBC, LFT and blood pressure at baseline Regular 3- monthly FBC, LFT and ophthalmology during therapy (34, 44, 45)	£19,169	\$82,043	2 nd line in HARRMS, or 1 st line in RESMS
Teriflunomide (Aubagio)	Daily oral tablet (14mg)	TOWER – 36.3% reduction in ARR compared to placebo, reduced risk of SAD by 31.5% (46) TEMPO – Reduced ARR by 31.5% compared to placebo, Reduced risk of SAD by 29.8% (47) TENORE – No difference in ARR compared to s.c. IFNβ1a (48)	Teriflunomide is thought to inhibit proliferation of activated T cells by inhibiting pyrimidine synthesis by the enzyme DHODH (dihydroorotate dehydrogenase) (49).	Elevated liver enzymes, Hair thinning, Nausea, Diarrhoea, Hypertension, Peripheral neuropathy (34, 46)	Baseline FBC, LFT, pregnancy test, TB, blood pressure First 6 months of therapy: monthly FBC, LFT After 6 months: FBC and LFT every 6 months (34)	£13,529.	\$76,612	RRMS
Dimethyl fumarate (Tecfidera)	120 mg twice daily oral	DEFINE –	Dimethyl fumarate has anti-inflammatory effects, by affecting	Nausea, Diarrhoea, Flushing (59),	Baseline FBC and LFT (34)	£17,898	\$82,977	RRMS

Pre-submission

	tablet; increased to 240 mg twice daily after 7 days	53% reduction in ARR compared to placebo, 38% reduction in risk of SAD (50) CONFIRM – 44% reduction in ARR compared to placebo, Significant reduction in ARR compared to GA, No reduction in SAD (51)	cytokine release and immune cell differentiation (52-56). It may also have neuroprotective antioxidant effects via Nrf2 (nuclear (erythroid-derived 2) related factor) pathways. In vitro studies showed that dimethyl fumarate protected neurones and glial cells against oxidative stress (57), and that this effect was lost in Nrf2 deficient cells (58). In vivo studies in EAE mice showed raised levels of Nrf2 and improved neuronal survival which was not seen in Nrf2 deficient mice (57).	Rarely PML (60)	Baseline MRI (60) Monitoring of white cell count is necessary as cases of PML have been reported in association with lymphopenia (61, 62)			
Natalizumab (Tysabri)	300mg i.v. infusion every 4 weeks.	AFFIRM – 68% reduction in ARR compared to placebo, and a 42% reduction in risk of SAD (63)	Natalizumab is a monoclonal antibody directed against $\alpha 4\beta 1$ integrin, an adhesion molecule involved in the extravasation of leukocytes across the BBB.	Mild infusion associated symptoms – headache and fever, Hypersensitivity reaction requiring discontinuation of infusion (34), Cases of hepatic injury have been reported (37) The risk of PML after two years of therapy is 1 in 100 in JCV seropositive patients who have received prior immunosuppressive treatment (64), Neutralising antibodies causing delayed hypersensitivity and reduced efficacy (65)	MRI at baseline, then every 3-12 months during therapy based on risk followed by one 3-6 months post-therapy (66) JCV serology every 6 months(66) Consider anti-natalizumab antibody testing after 6 months (65)	£14,730 annually	\$78,214	2 nd line for HARRMS or 1 st line for RESMS
Alemtuzumab (Lemtrada)	First annual course is 60mg via i.v. infusion over 5 days. Second annual course is 36mg via i.v. infusion over 3 days.	CARE MS I – (Alemtuzumab as 1 st line therapy) 54.9% reduction in ARR and a 30% reduction in risk of SAD compared to 44µg s.c. IFNβ1a (67) CARE MS II – (Alemtuzumab as 2 nd line therapy) 49.4% reduction in ARR and a 42% reduction in risk of SAD compared with 44µg s.c. IFNβ1a (68)	Alemtuzumab is a mAb targeting CD52, a surface marker on lymphocytes and monocytes. Alemtuzumab treatment depletes lymphocytes and monocytes which subsequently replete over a period of time, at different rates – CD4+ T cells take more than 5 years to reach pre-treatment levels (69). B cells replete over three months and then rise beyond pre-treatment levels, and this is thought to contribute to the secondary autoimmune complications which frequently follow alemtuzumab therapy (69, 70).	Infusion related reactions (90%) - may warrant pre-treatment with steroid, antihistamine and paracetamol (71), Infections, Secondary autoimmune conditions develop in up to 50% of patients after 2 years of therapy (70), Neutralizing antibodies occur in up to 80% of cases after second annual course (68)	Monthly blood tests including FBC, TFT, LFT Consider testing for neutralizing antibodies (71)	£56,360 for full two year course	\$103,749	Active RRMS

Pre-submission

Ocrelizumab (Ocrevus)	600mg i.v. infusion every 6 months	OPERA I and II – 46% -47% reduction in ARR and 40% reduction in risk of SAD compared to IFNb (72) ORATORIO – 24% and 25% reduction in risk of SAD at 12 and 24 weeks respectively compared to placebo (73)	Ocrelizumab targets CD20, a B cell progenitor antigen, and thereby depletes B cells (12)	Infusion-related reactions, Infections (72)	FBC (74)	£19,160 (£4790 per 300mg)	\$65,000	RRMS PPMS
Cladribine (Mavenclad)	1 or 2 10mg oral tablet once per day for four or five days of 2 treatment weeks each year for a total dose of 3.5mg/kg over 2 years	CLARITY – 57.6% and 54.5% reduction in ARR compared to placebo for the 3.5mg/kg and the 5.25mg/kg groups respectively. 33% and 31% reduction in risk of SAD for the 3.5mg/kg and 5.25mg/kg groups respectively compared to placebo (75)	Cladribine is an analogue of deoxyadenosine, which is metabolised to an active compound which causes cell death by incorporating into DNA and inhibiting DNA repair. It rapidly depletes T and B cells (76)	lymphopenia, infections – notably herpes zoster infections (77)	FBC Screen for HIV, TB and Hepatitis before commencing treatment (78)	£2,047.24 per 10mg tablet: Total cost of £50,157 for the 2 year course for a 70kg person	N/A	HARRMS

WAC = wholesale acquisition cost; s.c. = subcutaneous; i.m. = intramuscular; i.v. = intravenous; FBC = full blood count; LFT = liver function tests; TFT = thyroid function tests; PML = progressive multifocal leukoencephalopathy; JCV = John Cunningham virus; VZV = varicella zoster virus; TB = tuberculosis; RRMS = relapsing-remitting multiple sclerosis; HARRMS = highly active relapsing remitting multiple sclerosis; RESMS = rapidly evolving severe multiple sclerosis; ARR = Annualised Relapse Rate, Relapses are usually defined as worsening of neurological signs and symptoms lasting at least 24 or 48 hours, not associated with fever or infection, on a background of at least 30 days clinical stability; EDSS = (Kurtzke) Expanded Disability Status Scale; SAD = Sustained Accumulation of Disability, usually defined as 1 or more point increase in EDSS sustained for at least 3 or 6 months, or 0.5 point increase if baseline EDSS is over 5.5.

*= These are list prices, actual prices may be lower

First-line Injectable Therapies

These were the first DMTs developed for MS, and are still the most commonly used. They are favoured as they have a good safety profile (see Table 1) and have been in clinical use the longest. However, they are less effective than most newer DMTs (Table 1) (22, 23, 35, 38). Of course, there are issues in simply comparing outcome figures across different trials – with varying methods, patient characteristics, outcome definitions, trial lengths performed, safety profiles and trials performed at different times over recent history. A recent meta-analysis showed that although IFN and GA were generally among the least effective DMTs when looking at Annualised Relapse Rate (ARR) and 3-month Sustained Accumulation of Disability (SAD), IFNb1b came out as the most effective in reducing the risk of 6-month SAD, and pegylated IFNb1a was third on the list for reducing 6-month SAD (79). Still, cost-effectiveness models consistently show fewer QALYs gained with IFN and GA compared to other DMTs, even though these models generally take into account both ARR and SAD (80-82).

These drugs were initially deemed by the National Institute for Clinical Excellence (NICE), within the UK, to not meet their cost-benefit threshold, but were provided by the NHS under the risk-sharing scheme (RSS), in which the drug would be approved by NICE for use in the NHS if it was found to be cost-effective in a 20-year model based off of 10 years of follow-up data, and if not the price of the drugs for these patients would be discounted by the manufacturer (83). The outcome of the re-evaluation by NICE in early 2018 further concluded glatiramer acetate and all the beta interferons

except Betaferon are recommended for use in people with RRMS but only if provided with a discount under commercial agreement with suppliers (84). Betaferon was not considered to be cost-effective (84).

Oral Therapies

Current oral therapies for MS include teriflunomide, fingolimod, dimethyl fumarate and oral cladribine prodrug (Table 1). Although these oral therapies have the advantage of not being injected, they do still require regular blood test monitoring. In terms of effectiveness, Teriflunomide is as effective as IFN and GA, whilst dimethyl fumarate and fingolimod are more effective (79). However, these are still not as effective as the second-line injectable therapies natalizumab and alemtuzumab (79). The annual cost of these drugs varies from £13,500-£25,000, making them relatively expensive compared to IFN and GA. Oral cladribine is arguably the most effective of the oral agents for MS (77). In a recent cost effective analysis (CEA), cladribine was found to be more cost-effective than alemtuzumab and natalizumab (85). This may in part account for the relatively rapid approval of oral cladribine by NICE and availability to the NHS shortly following approval by the European Medicines agency. In terms of safety, these oral DMTs are all relatively safe compared to DMTs such as natalizumab and alemtuzumab (see Table 1).

Second-line Therapies

Natalizumab was the first monoclonal antibody licensed for use in MS. It is one of the most effective therapies available (79). At £14,730 annually, it is one of the less expensive DMTs for MS. However, this is not taking into account the cost for any additional monitoring required. Indeed, for one CEA the price per QALY gained went from 38,000 EUR (£32,000) when looking only at direct healthcare costs, to 59,000 EUR (£50,000) when taking into account the extra monitoring required due to the risk of progressive multifocal leukoencephalopathy (PML) (80).

PML is a progressive inflammatory condition in the brain caused by the John Cunningham virus (JCV). It can be associated with immunosuppression and a low white cell count (86). Currently, the only treatment is to terminate immunosuppressive therapy in order to minimise disease progression. It has a high mortality rate, approximately 20-25% at 6 months (87, 88). This is a rare occurrence with fingolimod and dimethyl fumarate therapy (44, 89). Though no cases have yet been reported with ocrelizumab therapy (unless following natalizumab therapy (90)), PML has been known to occur with anti-CD20 antibodies (91). The greatest risk of PML reported so far occurs with natalizumab therapy. The risk of PML after two years of therapy is about 1 in 100 in JCV seropositive patients who have received prior immunosuppressive treatment (64). Therefore natalizumab therapy requires MRI monitoring and JC virus serology status is monitored at 6-monthly intervals whilst on therapy (92).

Alemtuzumab is another monoclonal antibody that is currently licensed for RRMS. According to a recent meta-analysis, it is the most effective treatment for RRMS when looking at outcomes such as ARR improvement and preventing disability progression (79). However, due to the fact that disease progression is not always defined by the same criteria in different studies, and people in the pivotal alemtuzumab trials had a shorter disease course and therefore perhaps more neurological reserve to respond to therapy, the evidence for efficacy in preventing disease progression is less reliable (79). Significant monitoring is needed as the risk for secondary autoimmunity is as high as 50% of those treated with two annual courses (70). The cost of these potential complications is high as they include

thyroid autoimmunity, ITP (immune thrombocytopaenic purpura) and renal anti-glomerular basement membrane disease (70). These are costly to manage and can lead to severe disability and death (70). Furthermore, neutralizing antibodies occur in up to 30% of patients following the first course (68). This incidence increases to 80% after the second annual course (68). In many cases neutralizing antibody levels decrease over the year between treatment courses, rendering them non-problematic, but in some cases they persist causing treatments to be ineffective (68). Repeated courses past two years may not be necessary though as effects can last up to 6 years and possibly beyond (93-96).

Two phase III trials were conducted with ocrelizumab for RRMS, and one for PPMS (72, 73). Its efficacy data in these pivotal trials was comparable to current most effective DMTs. SAEs occurred at a similar rate as with IFN treatment or placebo, making it relatively safe compared to most other DMTs (72, 73). However, development of ocrelizumab was halted for rheumatoid arthritis and lupus due to deaths from infection (97, 98). It remains to be seen what the long-term safety profile of this drug will be like. Currently, it is administered every 6 months, however the phase II extension study suggests that like alemtuzumab and oral cladribine, that ocrelizumab may have induction therapy potential with long-term benefit from short courses of therapy (12, 13). Although cost-benefit value of ocrelizumab is accepted, this agent has yet to be seen as sufficiently cost-effective for treatment of PPMS. Although the manufacturers had offered supply to the NHS at a lower price than for use in relapsing MS variants, NICE has yet to entertain this model (99).

Haematopoietic Stem Cell Transplantation

A recent study showed total suppression of CNS inflammatory processes with autologous haematopoietic stem cell transplant (aHSCT), evidenced by a complete lack of relapses in any patients during a 7-year follow up period (100). Disability progression was halted in 70% of patients (100). This is similar to other studies looking at aHSCT for MS, but the total suppression of CNS immunity had not been seen before (101-103). Effects seen have been dramatic, but the procedure is currently reserved for last resort cases due to the high associated risk of mortality at around 5% (104), although this can be minimised in specialist centres to 1-2% (105). If safety improvements can be made and high efficacy results continue to be seen, then the procedure may become more commonplace in the future.

Cost-Effectiveness Analyses

The CEAs for MS drugs use models to predict the long-term cost and benefit of drugs based on published trial data. There are various issues with this. CEAs can take on the societal or the third-party payers perspective. The third party payers perspective includes only the cost of the drug and the cost to healthcare providers, whereas societal perspective will include informal care costs and indirect costs, such as the cost of unemployment. Thus, changing the perspective will impact the cost-effectiveness of the drug. A societal perspective is arguably more useful and realistic, but also less reliable than the payers perspective. Indeed, methods for estimating costs to society can vary widely between CEAs (106).

Changing the length of time of the model can affect the outcome of CEAs (107), as can using different sources for utility values (108). Some models do not account for non-drug care costs or the costs of complications and side effects of treatment. Many CEAs fail to transparently describe and justify their methods of cost estimation and the limitations of their studies, and sponsored CEAs tend to favour

the drug of the sponsor (106, 107). Furthermore, as the payer threshold per QALY is known, health economic data may be adapted to fit this threshold.

A CEA is only as good as the original trial data being used, therefore models using data from longer and larger trials with similar study populations are preferable. Head-to-head comparative randomized controlled trials (RCTs) are ideal but there is a near total dearth of RCTs comparing any DMT with any other DMT other than IFN or GA (79). Yet there are a number of cost-effectiveness models comparing various DMTs with each other, using data from separate trials or from historical comparator populations (106, 109). However comparisons are emerging following comparisons of real life usage using registries such as NARCOMS (North American) and MSBase (110).

The RCTs need to minimise the heterogeneity of outcome measures used. Defining SAD by 3 months instead of 6 may overestimate the rate of disability progression. Similarly, defining a relapse as lasting 24 verses 48 h can impact the results. Furthermore, as treatments become more effective these outcome measures become less relevant. There has been a move towards inclusion of other end points, notably that of the proportion of patients with no evidence of disease activity (NEDA). NEDA has been defined as “absence of new or enlarging T2 lesions or T1 gadolinium-enhancing lesions on MRI and no sustained EDSS score progression or clinical relapse” (111). In light of the improving efficacy of therapies, NEDA has become an important end-point. However, current definitions of NEDA may not pick up slight changes in disability progression (112). Furthermore, none of these outcome measures take into account the impact of MS and treatments on things like mental health, working status and quality of life, all of which may be more pertinent to look at as treatments are becoming more effective at suppressing disease activity as measured by relapses, MRI lesions and EDSS progression. As such NEDA evolves as new outcome measures form part of the definition such as brain atrophy and neurofilament levels (113).

A problem with long-term CEAs is that they cannot be challenged in a reasonable timeframe. There has been a call recently for manufacturers to support claims from CEAs with a protocol for empirical evaluation over a reasonable short-term timeframe, so that decision makers can confirm decisions on whether treatments shall be made available in their healthcare system (114). Long-term CEAs are arguably more useful to estimate the overall cost-benefit of a drug, especially when looking at whether more expensive but effective drugs can be a more cost-effective option in the context of a lifetime. However, these predictions cannot be challenged or tested, and as such cannot provide any real evidence of cost-effectiveness. Using cost per relapse avoided or year of NEDA instead of QALYs gained provides outcomes that can be verified in short-term studies. (83).

For the reasons outlined above, it is difficult to draw any conclusions from CEAs about which DMT is most cost effective.

Front-Loading of Costs

Recently, evidence is pointing towards the fact that front-loading of treatment costs in MS may be more cost-effective than paying for longer-term treatments, which are cheaper. It seems that drugs that are better at depleting memory B cells are more effective, and that induction of depletion may be all that is needed, without sustained treatment (12, 76). Furthermore, there is evidence to suggest that treating earlier in the disease course is more effective (115).

Currently, some of the more expensive and effective DMTs are showing promise as induction therapies with repeated courses after 1-2 years not being necessary, thus lowering the cost per QALY. Some of the best efficacy results have been seen with drugs such as alemtuzumab, ocrelizumab and oral cladribine, and there is evidence for all of these that treatment beyond the first couple of years may not be necessary (72, 93-96, 116). This means that though these are among the most expensive DMTs for MS, the cost over a lifetime may actually be less than some of the cheaper DMTs that require longer periods of treatment (85, 107, 117-119). Furthermore, treating aggressively early on in the disease to achieve NEDA may reduce hospital visits and admissions, further reducing the cost to healthcare systems and society (120). Indeed, one retrospective cohort study looking at the relationship between healthcare costs and disease progression in Italy found that higher care spending was associated with reduced progression over a 14 year time period (121). Long-term benefit from short treatment cycles means that the DMT are only present for short period and therefore may reduce problems of drug-drug interactions with the polypharmacopia of agents that will be needed to deal with symptomatic issues and neurorepair and protection. However, problems with these DMTs include the fact that they are new and long-term safety has not yet been determined. Furthermore, long-term efficacy has not been confirmed beyond a few years.

Conclusion

MS is a progressively disabling disease that has a significant impact on quality of life. This has implications for the individual with MS, for their families and for Society as people with MS are more likely to be unemployed and to require informal care from friends and family. This is in addition to the formal healthcare cost of managing the disease and the disability caused by it. Clearly there is a great need for treatments that can slow the progression of the disease and prevent disability.

Current first-line therapies are relatively safe and cheap but less effective compared to second-line therapies (79). There may be something to be said for using more expensive therapies that are also more effective to reduce the long-term risk of disability and to reduce long-term costs. Currently it is difficult to say what the long term outcomes of different therapies may be as there is not enough long-term follow-up data. Cost-effectiveness models have aimed to compare DMTs with each other over long-term time horizons, but their results are often difficult to interpret or compare across analyses. Currently, the drugs with the most benefit tend to be those that carry the most risk and highest price tag. It may be worth using more aggressive measures earlier on as front-loading the cost means the cost of long-term disability progression is reduced. Further head-to-head RCTs, RCT follow-up studies and CEAs are needed to answer these questions, and these need to take on board the recommendations that have been made for improving their quality. As patents expire notably on chemical small molecules that have low costs, the treatment landscape and cost-effectiveness requirements may change (122).

REFERENCES

1. Compston A, Coles A. Multiple sclerosis. *Lancet*. 2008;372(9648):1502-17.
2. Alonso A, Hernan MA. Temporal trends in the incidence of multiple sclerosis: a systematic review. *Neurology*. 2008;71(2):129-35.
3. Confavreux C, Aimard G, Devic M. Course and prognosis of multiple sclerosis assessed by the computerized data processing of 349 patients. *Brain*. 1980;103(2):281-300.

4. McDonald WI, Compston A, Edan G, Goodkin D, Hartung H-P, Lublin FD, et al. Recommended diagnostic criteria for multiple sclerosis: Guidelines from the international panel on the diagnosis of multiple sclerosis. *Annals of Neurology*. 2001;50(1):121-7.
5. Polman CH, Reingold SC, Banwell B, Clanet M, Cohen JA, Filippi M, et al. Diagnostic criteria for multiple sclerosis: 2010 Revisions to the McDonald criteria. *Annals of Neurology*. 2011;69(2):292-302.
6. Compston A, Coles A. Multiple sclerosis. *Lancet*. 2002;359(9313):1221-31.
7. de Vos AF, van Meurs M, Brok HP, Boven LA, Hintzen RQ, van der Valk P, et al. Transfer of central nervous system autoantigens and presentation in secondary lymphoid organs. *J Immunol*. 2002;169(10):5415-23.
8. van Zwam M, Huizinga R, Melief MJ, Wierenga-Wolf AF, van Meurs M, Voerman JS, et al. Brain antigens in functionally distinct antigen-presenting cell populations in cervical lymph nodes in MS and EAE. *J Mol Med (Berl)*. 2009;87(3):273-86.
9. Viglietta V, Baecher-Allan C, Weiner HL, Hafler DA. Loss of functional suppression by CD4+CD25+ regulatory T cells in patients with multiple sclerosis. *J Exp Med*. 2004;199(7):971-9.
10. Bartholomaeus I, Kawakami N, Odoardi F, Schlager C, Miljkovic D, Ellwart JW, et al. Effector T cell interactions with meningeal vascular structures in nascent autoimmune CNS lesions. *Nature*. 2009;462(7269):94-8.
11. Magliozzi R, Howell O, Vora A, Serafini B, Nicholas R, Puopolo M, et al. Meningeal B-cell follicles in secondary progressive multiple sclerosis associate with early onset of disease and severe cortical pathology. *Brain*. 2007;130(Pt 4):1089-104.
12. Baker D, Marta M, Pryce G, Giovannoni G, Schmierer K. Memory B Cells are Major Targets for Effective Immunotherapy in Relapsing Multiple Sclerosis. *EBioMedicine*. 2017;16:41-50.
13. Baker D, Pryce G, Amor S, Giovannoni G, Schmierer K. Learning from other autoimmunities to understand targeting of B cells to control multiple sclerosis. *Brain*. 2018.
14. Orme M, Kerrigan J, Tyas D, Russell N, Nixon R. The effect of disease, functional status, and relapses on the utility of people with multiple sclerosis in the UK. *Value Health*. 2007;10(1):54-60.
15. Murphy N, Confavreux C, Haas J, Konig N, Roullet E, Sailer M, et al. Quality of life in multiple sclerosis in France, Germany, and the United Kingdom. *J Neurol Neurosurg Psychiatry*. 1998;65(4):460-6.
16. Feuillet L, Reuter F, Audoin B, Malikova I, Barrau K, Cherif AA, et al. Early cognitive impairment in patients with clinically isolated syndrome suggestive of multiple sclerosis. *Mult Scler*. 2007;13(1):124-7.
17. Grytten Torkildsen N, Lie SA, Aarseth JH, Nyland H, Myhr KM. Survival and cause of death in multiple sclerosis: results from a 50-year follow-up in Western Norway. *Mult Scler*. 2008;14(9):1191-8.
18. Hoang H, Laursen B, Stenager EN, Stenager E. Psychiatric co-morbidity in multiple sclerosis: The risk of depression and anxiety before and after MS diagnosis. *Mult Scler*. 2016;22(3):347-53.
19. Pflieger CC, Flachs EM, Koch-Henriksen N. Social consequences of multiple sclerosis. Part 2. Divorce and separation: a historical prospective cohort study. *Mult Scler*. 2010;16(7):878-82.
20. Kobelt G, Thompson A, Berg J, Gannedahl M, Eriksson J. New insights into the burden and costs of multiple sclerosis in Europe. *Mult Scler*. 2017;1352458517694432.
21. Thompson A, Kobelt G, Berg J, Capsa D, Eriksson J, Miller D, et al. New insights into the burden and costs of multiple sclerosis in Europe: Results for the United Kingdom. <http://dxdoi.org/10.1177/1352458517708687>. 2017.
22. Randomised double-blind placebo-controlled study of interferon beta-1a in relapsing/remitting multiple sclerosis. PRISMS (Prevention of Relapses and Disability by Interferon beta-1a Subcutaneously in Multiple Sclerosis) Study Group. *Lancet*. 1998;352(9139):1498-504.
23. Jacobs LD, Cookfair DL, Rudick RA, Herndon RM, Richert JR, Salazar AM, et al. Intramuscular interferon beta-1a for disease progression in relapsing multiple sclerosis. The Multiple Sclerosis Collaborative Research Group (MSCRG). *Ann Neurol*. 1996;39(3):285-94.

24. Jiang H, Milo R, Swoveland P, Johnson KP, Panitch H, Dhib-Jalbut S. Interferon beta-1b reduces interferon gamma-induced antigen-presenting capacity of human glial and B cells. *J Neuroimmunol.* 1995;61(1):17-25.
25. Teleshova N, Bao W, Kivisakk P, Ozenci V, Mustafa M, Link H. Elevated CD40 ligand expressing blood T-cell levels in multiple sclerosis are reversed by interferon-beta treatment. *Scand J Immunol.* 2000;51(3):312-20.
26. Liu Z, Pelfrey CM, Cotleur A, Lee JC, Rudick RA. Immunomodulatory effects of interferon beta-1a in multiple sclerosis. *J Neuroimmunol.* 2001;112(1-2):153-62.
27. Saresella M, Marventano I, Longhi R, Lissoni F, Trabattoni D, Mendozzi L, et al. CD4+CD25+FoxP3+PD1- regulatory T cells in acute and stable relapsing-remitting multiple sclerosis and their modulation by therapy. *Faseb j.* 2008;22(10):3500-8.
28. Muraro PA, Liberati L, Bonanni L, Pantalone A, Caporale CM, Iarlori C, et al. Decreased integrin gene expression in patients with MS responding to interferon-beta treatment. *J Neuroimmunol.* 2004;150(1-2):123-31.
29. Muraro PA, Leist T, Bielekova B, McFarland HF. VLA-4/CD49d downregulated on primed T lymphocytes during interferon-beta therapy in multiple sclerosis. *J Neuroimmunol.* 2000;111(1-2):186-94.
30. Hunt D, Kavanagh D, Drummond I, Weller B, Bellamy C, Overell J, et al. Thrombotic microangiopathy associated with interferon beta. *N Engl J Med.* 2014;370(13):1270-1.
31. Vosoughi R, Freedman MS. Therapy of MS. *Clin Neurol Neurosurg.* 2010;112(5):365-85.
32. Schmidt S, Hertfelder HJ, von Spiegel T, Hering R, Harzheim M, Lassmann H, et al. Lethal capillary leak syndrome after a single administration of interferon beta-1b. *Neurology.* 1999;53(1):220-2.
33. Neurologists AoB. Revised (2009) Association of British Neurologists' guidelines for prescribing in multiple sclerosis 2009 [Available from: [http://www.theabn.org/media/docs/ABN%20publications/ABN_MS_Guidelines_2009_Final\(1\).pdf](http://www.theabn.org/media/docs/ABN%20publications/ABN_MS_Guidelines_2009_Final(1).pdf).
34. Cross AH, Naismith RT. Established and novel disease-modifying treatments in multiple sclerosis. *J Intern Med.* 2014;275(4):350-63.
35. Paty DW, Li DK. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. II. MRI analysis results of a multicenter, randomized, double-blind, placebo-controlled trial. UBC MS/MRI Study Group and the IFNB Multiple Sclerosis Study Group. *Neurology.* 1993;43(4):662-7.
36. Calabresi PA, Kieseier BC, Arnold DL, Balcer LJ, Boyko A, Pelletier J, et al. Pegylated interferon beta-1a for relapsing-remitting multiple sclerosis (ADVANCE): a randomised, phase 3, double-blind study. *Lancet Neurol.* 2014;13(7):657-65.
37. EMA. Plegidry - Summary of product characteristics. European Medicines Agency; 2014.
38. Johnson KP, Brooks BR, Cohen JA, Ford CC, Goldstein J, Lisak RP, et al. Copolymer 1 reduces relapse rate and improves disability in relapsing-remitting multiple sclerosis: results of a phase III multicenter, double-blind placebo-controlled trial. The Copolymer 1 Multiple Sclerosis Study Group. *Neurology.* 1995;45(7):1268-76.
39. Miller A, Shapiro S, Gershtein R, Kinarty A, Rawashdeh H, Honigman S, et al. Treatment of multiple sclerosis with copolymer-1 (Copaxone): implicating mechanisms of Th1 to Th2/Th3 immune-deviation. *J Neuroimmunol.* 1998;92(1-2):113-21.
40. Racke MK, Martin R, McFarland H, Fritz RB. Copolymer-1-induced inhibition of antigen-specific T cell activation: interference with antigen presentation. *J Neuroimmunol.* 1992;37(1-2):75-84.
41. Calabresi PA, Radue EW, Goodin D, Jeffery D, Rammohan KW, Reder AT, et al. Safety and efficacy of fingolimod in patients with relapsing-remitting multiple sclerosis (FREEDOMS II): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Neurol.* 2014;13(6):545-56.
42. Cohen JA, Barkhof F, Comi G, Hartung HP, Khatri BO, Montalban X, et al. Oral fingolimod or intramuscular interferon for relapsing multiple sclerosis. *N Engl J Med.* 2010;362(5):402-15.
43. Mehling M, Kappos L, Derfuss T. Fingolimod for multiple sclerosis: mechanism of action, clinical outcomes, and future directions. *Curr Neurol Neurosci Rep.* 2011;11(5):492-7.

44. New recommendations to minimise risks of the rare brain infection PML and a type of skin cancer with Gilenya [press release]. EMA2015.
45. Rommer PS, Zettl UK, Kieseier B, Hartung HP, Menge T, Frohman E, et al. Requirement for safety monitoring for approved multiple sclerosis therapies: an overview. *Clin Exp Immunol*. 2014;175(3):397-407.
46. Confavreux C, O'Connor P, Comi G, Freedman MS, Miller AE, Olsson TP, et al. Oral teriflunomide for patients with relapsing multiple sclerosis (TOWER): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Neurol*. 2014;13(3):247-56.
47. O'Connor P, Wolinsky JS, Confavreux C, Comi G, Kappos L, Olsson TP, et al. Randomized trial of oral teriflunomide for relapsing multiple sclerosis. *N Engl J Med*. 2011;365(14):1293-303.
48. Vermersch P, Czlankowska A, Grimaldi LM, Confavreux C, Comi G, Kappos L, et al. Teriflunomide versus subcutaneous interferon beta-1a in patients with relapsing multiple sclerosis: a randomised, controlled phase 3 trial. *Mult Scler*. 2014;20(6):705-16.
49. Bar-Or A, Pachner A, Menguy-Vacheron F, Kaplan J, Wiendl H. Teriflunomide and its mechanism of action in multiple sclerosis. *Drugs*. 2014;74(6):659-74.
50. Gold R, Kappos L, Arnold DL, Bar-Or A, Giovannoni G, Selmaj K, et al. Placebo-controlled phase 3 study of oral BG-12 for relapsing multiple sclerosis. *N Engl J Med*. 2012;367(12):1098-107.
51. Fox RJ, Miller DH, Phillips JT, Hutchinson M, Havrdova E, Kita M, et al. Placebo-controlled phase 3 study of oral BG-12 or glatiramer in multiple sclerosis. *N Engl J Med*. 2012;367(12):1087-97.
52. Ghoreschi K, Bruck J, Kellerer C, Deng C, Peng H, Rothfuss O, et al. Fumarates improve psoriasis and multiple sclerosis by inducing type II dendritic cells. *J Exp Med*. 2011;208(11):2291-303.
53. Litjens NH, Rademaker M, Ravensbergen B, Rea D, van der Plas MJ, Thio B, et al. Monomethylfumarate affects polarization of monocyte-derived dendritic cells resulting in down-regulated Th1 lymphocyte responses. *Eur J Immunol*. 2004;34(2):565-75.
54. de Jong R, Bezemer AC, Zomerdijsk TP, van de Pouw-Kraan T, Ottenhoff TH, Nibbering PH. Selective stimulation of T helper 2 cytokine responses by the anti-psoriasis agent monomethylfumarate. *Eur J Immunol*. 1996;26(9):2067-74.
55. Asadullah K, Schmid H, Friedrich M, Randow F, Volk HD, Sterry W, et al. Influence of monomethylfumarate on monocytic cytokine formation--explanation for adverse and therapeutic effects in psoriasis? *Arch Dermatol Res*. 1997;289(11):623-30.
56. Zhu K, Mrowietz U. Inhibition of dendritic cell differentiation by fumaric acid esters. *J Invest Dermatol*. 2001;116(2):203-8.
57. Linker RA, Lee DH, Ryan S, van Dam AM, Conrad R, Bista P, et al. Fumaric acid esters exert neuroprotective effects in neuroinflammation via activation of the Nrf2 antioxidant pathway. *Brain*. 2011;134(Pt 3):678-92.
58. Scannevin RH, Chollate S, Jung MY, Shackett M, Patel H, Bista P, et al. Fumarates promote cytoprotection of central nervous system cells against oxidative stress via the nuclear factor (erythroid-derived 2)-like 2 pathway. *J Pharmacol Exp Ther*. 2012;341(1):274-84.
59. Gold R, Giovannoni G, Phillips JT, Fox RJ, Zhang A, Meltzer L, et al. Efficacy and safety of delayed-release dimethyl fumarate in patients newly diagnosed with relapsing-remitting multiple sclerosis (RRMS). *Mult Scler*. 2015;21(1):57-66.
60. Updated recommendations to minimise the risk of the rare brain infection PML with Tecfidera [press release]. EMA2015.
61. Salmen A, Gold R. Mode of action and clinical studies with fumarates in multiple sclerosis. *Exp Neurol*. 2014;262 Pt A:52-6.
62. Bomprezzi R. Dimethyl fumarate in the treatment of relapsing-remitting multiple sclerosis: an overview. *Ther Adv Neurol Disord*. 2015;8(1):20-30.
63. Polman CH, O'Connor PW, Havrdova E, Hutchinson M, Kappos L, Miller DH, et al. A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *N Engl J Med*. 2006;354(9):899-910.

64. Bloomgren G, Richman S, Hotermans C, Subramanyam M, Goelz S, Natarajan A, et al. Risk of Natalizumab-Associated Progressive Multifocal Leukoencephalopathy. <http://dxdoiorg/101056/NEJMoa1107829>. 2012.
65. Calabresi PA, Giovannoni G, Confavreux C, Galetta SL, Havrdova E, Hutchinson M, et al. The incidence and significance of anti-natalizumab antibodies: results from AFFIRM and SENTINEL. *Neurology*. 2007;69(14):1391-403.
66. McGuigan C, Craner M, Guadagno J, Kapoor R, Mazibrada G, Molyneux P, et al. Stratification and monitoring of natalizumab-associated progressive multifocal leukoencephalopathy risk: recommendations from an expert group. 2015.
67. Cohen JA, Coles AJ, Arnold DL, Confavreux C, Fox EJ, Hartung HP, et al. Alemtuzumab versus interferon beta 1a as first-line treatment for patients with relapsing-remitting multiple sclerosis: a randomised controlled phase 3 trial. *Lancet*. 2012;380(9856):1819-28.
68. Coles AJ, Twyman CL, Arnold DL, Cohen JA, Confavreux C, Fox EJ, et al. Alemtuzumab for patients with relapsing multiple sclerosis after disease-modifying therapy: a randomised controlled phase 3 trial. *The Lancet*. 380(9856):1829-39.
69. Coles AJ, Cox A, Le Page E, Jones J, Trip SA, Deans J, et al. The window of therapeutic opportunity in multiple sclerosis: evidence from monoclonal antibody therapy. *J Neurol*. 2006;253(1):98-108.
70. Willis MD, Harding KE, Pickersgill TP, Wardle M, Pearson OR, Scolding NJ, et al. Alemtuzumab for multiple sclerosis: Long term follow-up in a multi-centre cohort. *Mult Scler*. 2016;22(9):1215-23.
71. Coles AJ. Alemtuzumab therapy for multiple sclerosis. *Neurotherapeutics*. 2013;10(1):29-33.
72. Hauser SL, Bar-Or A, Comi G, Giovannoni G, Hartung HP, Hemmer B, et al. Ocrelizumab versus Interferon Beta-1a in Relapsing Multiple Sclerosis. *N Engl J Med*. 2017;376(3):221-34.
73. Montalban X, Hauser SL, Kappos L, Arnold DL, Bar-Or A, Comi G, et al. Ocrelizumab versus Placebo in Primary Progressive Multiple Sclerosis. *N Engl J Med*. 2017;376(3):209-20.
74. Agency EM. Ocrevus - Summary of Product Characteristics. EMA; 2018.
75. Giovannoni G, Comi G, Cook S, Rammohan K, Rieckmann P, Soelberg Sorensen P, et al. A placebo-controlled trial of oral cladribine for relapsing multiple sclerosis. *N Engl J Med*. 2010;362(5):416-26.
76. Ceronie B, Jacobs BM, Baker D, Dubuisson N, Mao Z, Ammoscato F, et al. Cladribine treatment of multiple sclerosis is associated with depletion of memory B cells. *J Neurol*. 2018.
77. Cook S, Vermersch P, Comi G, Giovannoni G, Rammohan K, Rieckmann P, et al. Safety and tolerability of cladribine tablets in multiple sclerosis: the CLARITY (CLAdRibine Tablets treating multiple sclerosis orally) study. *Mult Scler*. 2011;17(5):578-93.
78. Agency EM. Mavenclad - Summary of Product Characteristics. EMA; 2017.
79. Fogarty E, Schmitz S, Tubridy N, Walsh C, Barry M. Comparative efficacy of disease-modifying therapies for patients with relapsing remitting multiple sclerosis: Systematic review and network meta-analysis. *Mult Scler Relat Disord*. 2016;9:23-30.
80. Kobelt G, Berg J, Lindgren P, Jonsson B, Stawiarz L, Hillert J. Modeling the cost-effectiveness of a new treatment for MS (natalizumab) compared with current standard practice in Sweden. *Mult Scler*. 2008;14(5):679-90.
81. Earnshaw SR, Graham J, Oleen-Burkey M, Castelli-Haley J, Johnson K. Cost effectiveness of glatiramer acetate and natalizumab in relapsing-remitting multiple sclerosis. *Applied Health Economics and Health Policy*. 2009;7(2):91-108.
82. Gani R, Giovannoni G, Bates D, Kembell B, Hughes S, Kerrigan J. Cost-Effectiveness Analyses of Natalizumab (Tysabri®) Compared with Other Disease-Modifying Therapies for People with Highly Active Relapsing-Relapsing Multiple Sclerosis in the UK. *PharmacoEconomics*. 2008;26(7):617-27.
83. Duddy M, Palace J. The UK Risk-Sharing Scheme for interferon-beta and glatiramer acetate in multiple sclerosis. Outcome of the year-6 analysis. *Pract Neurol*. 2016;16(1):4-6.

84. NICE. NICE Technology appraisal guidance 527. Beta interferons and glatiramer acetate for treating multiple sclerosis. National Institute for Health and Care Excellence 2018 [Available from: <https://www.nice.org.uk/guidance/ta527>].
85. Hettle R, Harty G, Wong SL. Cost-effectiveness of cladribine tablets, alemtuzumab, and natalizumab in the treatment of relapsing-remitting multiple sclerosis with high disease activity in England. *J Med Econ*. 2018;1-11.
86. Pavlovic D, Patera AC, Nyberg F, Gerber M, Liu M. Progressive multifocal leukoencephalopathy: current treatment options and future perspectives. *Ther Adv Neurol Disord*. 2015;8(6):255-73.
87. Dong-Si T, Gheuens S, Gangadharan A, Wenten M, Philip J, McIninch J, et al. Predictors of survival and functional outcomes in natalizumab-associated progressive multifocal leukoencephalopathy. *J Neurovirol*. 2015;21(6):637-44.
88. Hellwig K, Gold R. Progressive multifocal leukoencephalopathy and natalizumab. *J Neurol*. 2011;258(11):1920-8.
89. Balak D, Hajdarbegovic E. More on PML in Patients Treated with Dimethyl Fumarate. *N Engl J Med*. 2016;374(3):295.
90. Kadish R, Robertson D, Sweeney M. Fatal leukoencephalopathy in a patient with multiple sclerosis following treatment with ocrelizumab (P5.353). 2018.
91. Sorensen PS, Blinkenberg M. The potential role for ocrelizumab in the treatment of multiple sclerosis: current evidence and future prospects. *Ther Adv Neurol Disord*. 2016. p. 44-52.
92. EMA confirms recommendations to minimise risk of brain infection PML with Tysabri [press release]. EMA2016.
93. Schippling S, Cohen J, Coles A, Compston DA, Fox E, Filippi M, et al. Alemtuzumab-Treated Patients with RRMS Demonstrate Durable Slowing of Brain Volume Loss Over 5 Years Despite Most Being Treatment-Free for 4 Years: CARE-MS I and II Extension Study (S51.001). *Neurology*. 2016;86(16 Supplement).
94. Limmroth V, Giovannoni G, Arnold D, Cohen J, Coles A, Fox E, et al. Treatment-Naive Patients with Active RRMS Demonstrate Durable Improvements in Relapse and Disability Following Treatment with Alemtuzumab: 5-Year Follow-Up of the CARE-MS I Study (S51.004). *Neurology*. 2016;86(16 Supplement).
95. Coles A, Arnold D, Cohen J, Fox E, Giovannoni G, Hartung H-P, et al. Patients with Active RRMS and an Inadequate Response to Prior Therapy Demonstrate Durable Improvements in Relapse and Disability Following Treatment with Alemtuzumab: 5-Year Follow-Up of the CARE-MS II Study (P3.022). *Neurology*. 2016;86(16 Supplement).
96. Arnold D, Cohen J, Coles A, Compston DA, Fox E, Hartung H-P, et al. Treatment-Naive Patients with Active RRMS Who Received Alemtuzumab Demonstrate Durable Suppression of New MRI Lesion Formation: 5-Year Follow-Up of the CARE-MS I Study (S51.002). *Neurology*. 2016;86(16 Supplement).
97. Emery P, Rigby W, Tak PP, Dörner T, Olech E, Martin C, et al. Safety with Ocrelizumab in Rheumatoid Arthritis: Results from the Ocrelizumab Phase III Program. *PLoS One*. 2014.
98. Mysler EF, Spindler AJ, Guzman R, Bijl M, Jayne D, Furie RA, et al. Efficacy and safety of ocrelizumab in active proliferative lupus nephritis: results from a randomized, double-blind, phase III study. *Arthritis Rheum*. 2013;65(9):2368-79.
99. NICE. Ocrelizumab for treating primary progressive multiple sclerosis [ID938]. National Institute for Health and Care Excellence 2018 [Available from: <https://www.nice.org.uk/guidance/indevelopment/gid-ta10153/documents>].
100. Atkins HL, Bowman M, Allan D, Anstee G, Arnold DL, Bar-Or A, et al. Immunoablation and autologous haemopoietic stem-cell transplantation for aggressive multiple sclerosis: a multicentre single-group phase 2 trial. *Lancet*. 2016;388(10044):576-85.
101. Nash RA, Hutton GJ, Racke MK, Popat U, Devine SM, Griffith LM, et al. High-dose immunosuppressive therapy and autologous hematopoietic cell transplantation for relapsing-remitting multiple sclerosis (HALT-MS): a 3-year interim report. *JAMA Neurol*. 2015;72(2):159-69.

102. Burman J, Iacobaeus E, Svenningsson A, Lycke J, Gunnarsson M, Nilsson P, et al. Autologous haematopoietic stem cell transplantation for aggressive multiple sclerosis: the Swedish experience. *J Neurol Neurosurg Psychiatry*. 2014;85(10):1116-21.
103. Burt RK, Balabanov R, Han X, Sharrack B, Morgan A, Quigley K, et al. Association of nonmyeloablative hematopoietic stem cell transplantation with neurological disability in patients with relapsing-remitting multiple sclerosis. *Jama*. 2015;313(3):275-84.
104. Snowden JA, Pearce RM, Lee J, Kirkland K, Gilleece M, Veys P, et al. Haematopoietic stem cell transplantation (HSCT) in severe autoimmune diseases: analysis of UK outcomes from the British Society of Blood and Marrow Transplantation (BSBMT) data registry 1997-2009. *Br J Haematol*. 2012;157(6):742-6.
105. Mancardi G, Saccardi R. Autologous haematopoietic stem-cell transplantation in multiple sclerosis. *Lancet Neurol*. 2008;7(7):626-36.
106. Guo S, Pelligra C, Saint-Laurent Thibault C, Hernandez L, Kansal A. Cost-effectiveness analyses in multiple sclerosis: a review of modelling approaches. *Pharmacoeconomics*. 2014;32(6):559-72.
107. Yamamoto D, Campbell JD. Cost-effectiveness of multiple sclerosis disease-modifying therapies: a systematic review of the literature. *Autoimmune Dis*. 2012;2012:784364.
108. Versteegh M. Impact on the Incremental Cost-Effectiveness Ratio of Using Alternatives to EQ-5D in a Markov Model for Multiple Sclerosis. *Pharmacoeconomics*. 2016;34(11):1133-44.
109. O'Day K, Meyer K, Stafkey-Mailey D, Watson C. Cost-effectiveness of natalizumab vs fingolimod for the treatment of relapsing-remitting multiple sclerosis: analyses in Sweden. *J Med Econ*. 2015;18(4):295-302.
110. Kalincik T, Brown JW, Robertson N, Willis M, Scolding N, Rice CM, et al. Treatment effectiveness of alemtuzumab compared with natalizumab, fingolimod, and interferon beta in relapsing-remitting multiple sclerosis: a cohort study. *Lancet Neurol*. 2017;16(4):271-81.
111. Rotstein DL, Healy BC, Malik MT, Chitnis T, Weiner HL. Evaluation of no evidence of disease activity in a 7-year longitudinal multiple sclerosis cohort. *JAMA Neurol*. 2015;72(2):152-8.
112. Stangel M, Penner IK, Kallmann BA, Lukas C, Kieseier BC. Towards the implementation of 'no evidence of disease activity' in multiple sclerosis treatment: the multiple sclerosis decision model. *Ther Adv Neurol Disord*. 2015;8(1):3-13.
113. Hakansson I, Tisell A, Cassel P, Blennow K, Zetterberg H, Lundberg P, et al. Neurofilament levels, disease activity and brain volume during follow-up in multiple sclerosis. *J Neuroinflammation*. 2018;15(1):209.
114. Schommer JC, Carlson AM, Rhee TG. Validating pharmaceutical product claims: questions a formulary committee should ask. *J Med Econ*. 2015;18(12):1000-6.
115. Tur C, Montalban X, Tintore M, Nos C, Rio J, Aymerich FX, et al. Interferon beta-1b for the treatment of primary progressive multiple sclerosis: five-year clinical trial follow-up. *Arch Neurol*. 2011;68(11):1421-7.
116. Comi G, Cook S, Rammohan K, Soelberg Sorensen P, Vermersch P, Adeniji AK, et al. Long-term effects of cladribine tablets on MRI activity outcomes in patients with relapsing-remitting multiple sclerosis: the CLARITY Extension study. *Ther Adv Neurol Disord*. 2018;11:1756285617753365.
117. Ontaneda D, Henry O. Cost Effectiveness of Multiple Sclerosis Therapeutic Strategies for Achieving No Evidence of Disease Activity (P2.193). 2016.
118. Smith A, Hashemi L, Ma I. Cost-Utility Analysis Of Alemtuzumab Versus Natalizumab For The Treatment Of Relapsing-Remitting Multiple Sclerosis: Us Payer Perspective. *Value in Health*. 2016;19(7).
119. Rog D, Salford Royal NHS Foundation Trust S, UK, Guo J, Sanofi C, MA, USA, Nucit A, Ividata L-P, France, et al. Alemtuzumab Is The Most Cost-Effective Option In Comparison To Available Therapies In The Treatment of Rrms From The UK NHS Perspective. *Value in Health*. 2017;20(9).
120. Cerqueira JJ, Compston DAS, Geraldes R, Rosa MM, Schmierer K, Thompson A, et al. Time matters in multiple sclerosis: can early treatment and long-term follow-up ensure everyone benefits from the latest advances in multiple sclerosis? *J Neurol Neurosurg Psychiatry*. 2018.

121. Moccia M, Palladino R, Lanzillo R, Carotenuto A, Russo CV, Triassi M, et al. Healthcare Costs for Treating Relapsing Multiple Sclerosis and the Risk of Progression: A Retrospective Italian Cohort Study from 2001 to 2015. *PLoS One*. 2017;12(1):e0169489.
122. Mao Z, Alvarez-Gonzalez C, De Trane S, Yildiz O, Albor C, Doctor G, et al. Cladribine: Off-label disease modification for people with multiple sclerosis in resource-poor settings? *Mult Scler J Exp Transl Clin*. 2018;4(2):2055217318783767.